

## **Anti-CCNB3** antibody



**Description** Unconjugated Rabbit polyclonal to CCNB3

Model STJ191900

**Host** Rabbit

**Reactivity** Human

**Applications** ELISA, WB

**Gene ID** 85417

Gene Symbol CCNB3

**Dilution range** WB 1:500-2000 ELISA 1:5000-20000

**Specificity** CCNB3 Polyclonal Antibody detects endogenous levels of protein.

**Tissue Specificity** Testis specific. In testis, it is expressed in developing germ cells, but not in

Leydig cells. Weakly or not expressed in other tissues.

**Purification** CCNB3 antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** G2/mitotic-specific cyclin-B3

Molecular Weight 153 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:18709OMIM:300456

Alternative Names G2/mitotic-specific cyclin-B3

**Function** Cyclins are positive regulatory subunits of the cyclin-dependent kinases

(CDKs), and thereby play an essential role in the control of the cell cycle, notably via their destruction during cell division. Its tissue specificity suggest

that it may be required during early meiotic prophase I.

Sequence and Domain Family The N-terminal destruction box (D-box) probably acts as a recognition signal

for degradation via the ubiquitin-proteasome pathway.

Cellular Localization Nucleus

**Post-translational** Ubiquitinated (Probable). Ubiquitination leads to its degradation during

**Modifications** anaphase entry, after degradation of CCNB1.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580 **T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com