

Anti-BAG2 antibody



Description	Unconjugated Rabbit polyclonal to BAG2
Model	STJ191961
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, WB
Gene ID	9532
Gene Symbol	BAG2
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	BAG2 Polyclonal Antibody detects endogenous levels of protein.
Purification	BAG2 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	BAG family molecular chaperone regulator 2 BAG-2 Bcl-2-associated athanogene 2
Molecular Weight	23 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:938OMIM:603882
Alternative Names	BAG family molecular chaperone regulator 2 BAG-2 Bcl-2-associated athanogene 2
Function	Co-chaperone for HSP70 and HSC70 chaperone proteins. Acts as a nucleotide-exchange factor (NEF) promoting the release of ADP from the HSP70 and HSC70 proteins thereby triggering client/substrate protein release.

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com