

Anti-TRIB2 antibody



| | |
|---------------------------|---|
| Description | Unconjugated Rabbit polyclonal to TRIB2 |
| Model | STJ191997 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Applications | ELISA, WB |
| Gene ID | 28951 |
| Gene Symbol | TRIB2 |
| Dilution range | WB 1:500-2000 ELISA 1:5000-20000 |
| Specificity | TRIB2 Polyclonal Antibody detects endogenous levels of protein. |
| Tissue Specificity | Highly expressed in peripheral blood leukocytes. |
| Purification | TRIB2 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Note | For Research Use Only (RUO). |
| Protein Name | Tribbles homolog 2 TRB-2 |
| Molecular Weight | 37 kDa |
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Isotype | IgG |
| Formulation | Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide. |

| | |
|-----------------------------------|--|
| Concentration | 1 mg/ml |
| Storage Instruction | Store at -20°C, and avoid repeat freeze-thaw cycles. |
| Database Links | HGNC:30809OMIM:609462 |
| Alternative Names | Tribbles homolog 2 TRB-2 |
| Function | Interacts with MAPK kinases and regulates activation of MAP kinases. Does not display kinase activity . |
| Sequence and Domain Family | The protein kinase domain is predicted to be catalytically inactive. |
| Cellular Localization | Cytoplasm Cytoplasm, cytoskeleton. May associate with the cytoskeleton. |

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com