

Anti-TR13C antibody



Description	Unconjugated Rabbit polyclonal to TR13C
Model	STJ192022
Host	Rabbit
Reactivity	Human
Applications	ELISA, WB
Gene ID	115650
Gene Symbol	TNFRSF13C
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	TR13C Polyclonal Antibody detects endogenous levels of protein.
Tissue Specificity	Highly expressed in spleen and lymph node, and in resting B-cells. Detected at lower levels in activated B-cells, resting CD4+ T-cells, in thymus and peripheral blood leukocytes.
Purification	TR13C antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor superfamily member 13C B-cell-activating factor receptor BAFF receptor BAFF-R BLyS receptor 3 CD antigen CD268
Molecular Weight	20 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:17755 OMIM:606269
Alternative Names	Tumor necrosis factor receptor superfamily member 13C B-cell-activating factor receptor BAFF receptor BAFF-R BLyS receptor 3 CD antigen CD268
Function	B-cell receptor specific for TNFSF13B/TALL1/BAFF/BLyS. Promotes the survival of mature B-cells and the B-cell response.
Cellular Localization	Membrane

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com