

## **Anti-CLC10** antibody



**Description** Unconjugated Rabbit polyclonal to CLC10

Model STJ192024

**Host** Rabbit

**Reactivity** Human, Mouse, Rat

**Applications** ELISA, WB

**Gene ID** 10462

Gene Symbol <u>CLEC10A</u>

**Dilution range** WB 1:500-2000 ELISA 1:5000-20000

**Specificity** CLC10 Polyclonal Antibody detects endogenous levels of protein.

**Purification** CLC10 antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

Protein Name C-type lectin domain family 10 member A C-type lectin superfamily member

14 Macrophage lectin 2 CD antigen CD301

**Molecular Weight** 34 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:16916OMIM:605999

Alternative Names C-type lectin domain family 10 member A C-type lectin superfamily member

14 Macrophage lectin 2 CD antigen CD301

**Function** Probable role in regulating adaptive and innate immune responses. Binds in a

calcium-dependent manner to terminal galactose and N-acetylgalactosamine units, linked to serine or threonine. These sugar moieties are known as Tn-Ag

and are expressed in a variety of carcinoma cells.

**Cellular Localization** Membrane