

## Anti-I27RA antibody



<b>Description</b>	Unconjugated Rabbit polyclonal to I27RA
--------------------	---

<b>Model</b>	STJ192032
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA, WB
<b>Gene ID</b>	<a href="#">9466</a>
<b>Gene Symbol</b>	<a href="#">IL27RA</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	I27RA Polyclonal Antibody detects endogenous levels of protein.
<b>Tissue Specificity</b>	Highly expressed in lymphoid tissues such as spleen, lymph nodes and peripheral blood leukocytes. Weakly expressed in other tissues examined including heart, brain, fetal and adult lung, liver, skeletal muscle, kidney, pancreas, prostate, testis, ovary, small intestine, kidney and colon. In the lymphoid system, higher level expression in CD4+ T-cell subsets than in CD8+ T-cell subsets. Also weaker expression in CD19+ B-cells and monocytes.
<b>Purification</b>	I27RA antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Interleukin-27 receptor subunit alpha IL-27 receptor subunit alpha IL-27R subunit alpha IL-27R-alpha IL-27RA Cytokine receptor WSX-1 Cytokine receptor-like 1 Type I T-cell cytokine receptor TCCR ZcytoR

<b>Molecular Weight</b>	69 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:17290</a> <a href="#">OMIM:605350</a>
<b>Alternative Names</b>	Interleukin-27 receptor subunit alpha IL-27 receptor subunit alpha IL-27R subunit alpha IL-27R-alpha IL-27RA Cytokine receptor WSX-1 Cytokine receptor-like 1 Type I T-cell cytokine receptor TCCR ZcytoR
<b>Function</b>	Receptor for IL27. Requires IL6ST/gp130 to mediate signal transduction in response to IL27. This signaling system acts through STAT3 and STAT1. Involved in the regulation of Th1-type immune responses. Also appears to be involved in innate defense mechanisms.
<b>Sequence and Domain Family</b>	The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.; The box 1 motif is required for JAK interaction and/or activation.
<b>Cellular Localization</b>	Membrane. Single-pass type I membrane protein.

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**W** <http://www.stjohnslabs.com/>

**T** +44 (0)208 223 3081

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)