

## Anti-N2DL3 antibody



**Description** Unconjugated Rabbit polyclonal to N2DL3

Model STJ192047

**Host** Rabbit

**Reactivity** Human

**Applications** ELISA, WB

**Gene ID** 79465

Gene Symbol <u>ULBP3</u>

**Dilution range** WB 1:500-2000 ELISA 1:5000-20000

**Specificity** N2DL3 Polyclonal Antibody detects endogenous levels of protein.

**Purification** N2DL3 antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

Protein Name UL16-binding protein 3 ALCAN-gamma NKG2D ligand 3 N2DL-3

NKG2DL3 Retinoic acid early transcript 1N

Molecular Weight 26 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:14895OMIM:605699</u>

Alternative Names UL16-binding protein 3 ALCAN-gamma NKG2D ligand 3 N2DL-3

NKG2DL3 Retinoic acid early transcript 1N

Function Ligand for the KLRK1/NKG2D receptor, together with at least ULBP1 and

ULBP2. ULBPs activate multiple signaling pathways in primary NK cells, resulting in the production of cytokines and chemokines. Binding of ULBPs ligands to KLRK1/NKG2D induces calcium mobilization and activation of the JAK2, STAT5, ERK and PI3K kinase/Akt signal transduction pathway. Has lower affinity for KLRK1/NKG2D compared to ULBP1 and ULBP2 and

induces weaker signaling responses than does ULBP2 or ULBP1.

Cellular Localization Cell membrane. Lipid-anchor, GPI-anchor.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580 **T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com