

## Anti-FLIP1 antibody

---



<b>Description</b>	Unconjugated Rabbit polyclonal to FLIP1
<b>Model</b>	STJ192087
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, WB
<b>Gene ID</b>	<a href="#">27145</a>
<b>Gene Symbol</b>	<a href="#">FILIP1</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	FLIP1 Polyclonal Antibody detects endogenous levels of protein.
<b>Tissue Specificity</b>	Moderately expressed in adult heart and brain. Weakly expressed in lung, skeletal muscle, ovary, testis, kidney, and fetal brain, and hardly detectable in liver, pancreas, spleen, and fetal liver. Within brain, moderate expression is found in amygdala and caudate nucleus.
<b>Purification</b>	FLIP1 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Filamin-A-interacting protein 1 FILIP
<b>Molecular Weight</b>	133 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated

<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:21015OMIM:607307</a>
<b>Alternative Names</b>	Filamin-A-interacting protein 1 FILIP
<b>Function</b>	By acting through a filamin-A/F-actin axis, it controls the start of neocortical cell migration from the ventricular zone. May be able to induce the degradation of filamin-A.
<b>Cellular Localization</b>	Cytoplasm, cytoskeleton

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)