

## Anti-PORIM antibody

---



<b>Description</b>	Unconjugated Rabbit polyclonal to PORIM
<b>Model</b>	STJ192279
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, WB
<b>Gene ID</b>	<a href="#">114908</a>
<b>Gene Symbol</b>	<a href="#">TMEM123</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	PORIM Polyclonal Antibody detects endogenous levels of protein.
<b>Tissue Specificity</b>	Ubiquitous. Not expressed in ovary. Expressed in keratinocytes.
<b>Purification</b>	PORIM antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Porimin Keratinocytes-associated transmembrane protein 3 KCT-3 Pro-oncosis receptor inducing membrane injury Transmembrane protein 123
<b>Molecular Weight</b>	22 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG

<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:30138</a> <a href="#">OMIM:606356</a>
<b>Alternative Names</b>	Porimin Keratinocytes-associated transmembrane protein 3 KCT-3 Pro-oncosis receptor inducing membrane injury Transmembrane protein 123
<b>Function</b>	Implicated in oncotic cell death, characterized by cell swelling, organelle swelling, vacuolization and increased membrane permeability.
<b>Cellular Localization</b>	Membrane

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580  
**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>  
**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)