

## **Anti-SARM1** antibody



**Description** Unconjugated Rabbit polyclonal to SARM1

Model STJ192592

**Host** Rabbit

**Reactivity** Human, Mouse

**Applications** ELISA, WB

**Immunogen** Synthesized peptide derived from human SARM1 protein.

**Immunogen Region** 281-330aa

**Gene ID** 23098

Gene Symbol SARM1

**Dilution range** WB 1:500-2000 ELISA 1:5000-20000

**Specificity** SARM1 Polyclonal Antibody detects endogenous levels of protein.

**Tissue Specificity** Predominantly expressed in brain, kidney and liver. Expressed at lower level

in placenta.

**Purification** SARM1 antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** Sterile alpha and TIR motif-containing protein 1 Sterile alpha and Armadillo

repeat protein Sterile alpha motif domain-containing protein 2 MyD88-5 SAM

domain-containing protein 2 Tir-1 homolog

Molecular Weight 79 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:17074OMIM:607732</u>

Alternative Names Sterile alpha and TIR motif-containing protein 1 Sterile alpha and Armadillo

repeat protein Sterile alpha motif domain-containing protein 2 MyD88-5 SAM

domain-containing protein 2 Tir-1 homolog

**Function** Negative regulator of MYD88- and TRIF-dependent toll-like receptor

signaling pathway which plays a pivotal role in activating axonal degeneration following injury. Promotes Wallerian degeneration an injury-induced axonal death pathway which involves degeneration of an axon distal to the injury site.

Can activate neuronal death in response to stress. Regulates dendritic arborization through the MAPK4-JNK pathway. Involved in innate immune response. Inhibits both TICAM1/TRIF- and MYD88-dependent activation of JUN/AP-1, TRIF-dependent activation of NF-kappa-B and IRF3, and the

phosphorylation of MAPK14/p38.

**Cellular Localization** Cytoplasm. Cell projection, axon Cell projection, dendrite Cell junction,

synapse Mitochondrion. Associated with microtubules.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com