

Anti-CAN1 antibody



Description	Unconjugated Rabbit polyclonal to CAN1
Model	STJ192952
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Gene ID	823
Gene Symbol	CAPN1
Dilution range	WB 1:500-2000 ELISA 1:5000-20000
Specificity	CAN1 Polyclonal Antibody detects endogenous levels of protein.
Tissue Specificity	Ubiquitous.
Purification	CAN1 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Calpain-1 catalytic subunit Calcium-activated neutral proteinase 1 CANP 1 Calpain mu-type Calpain-1 large subunit Cell proliferation-inducing gene 30 protein Micromolar-calpain muCANP
Molecular Weight	78 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:14760MIM:114220
Alternative Names	Calpain-1 catalytic subunit Calcium-activated neutral proteinase 1 CANP 1 Calpain mu-type Calpain-1 large subunit Cell proliferation-inducing gene 30 protein Micromolar-calpain muCANP
Function	Calcium-regulated non-lysosomal thiol-protease which catalyzes limited proteolysis of substrates involved in cytoskeletal remodeling and signal transduction.
Cellular Localization	Cytoplasm Cell membrane. Translocates to the plasma membrane upon Ca(2+) binding. In granular keratinocytes and in lower corneocytes, colocalizes with FLG and FLG2 .
Post-translational Modifications	Undergoes calcium-induced successive autoproteolytic cleavages that generate a membrane-bound 78 kDa active form and an intracellular 75 kDa active form. Calpastatin reduces with high efficiency the transition from 78 kDa to 75 kDa calpain forms.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com