

## Anti-COAA1 antibody

---



<b>Description</b>	Unconjugated Rabbit polyclonal to COAA1
<b>Model</b>	STJ192990
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA, WB
<b>Gene ID</b>	<a href="#">1300</a>
<b>Gene Symbol</b>	<a href="#">COL10A1</a>
<b>Dilution range</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Specificity</b>	COAA1 Polyclonal Antibody detects endogenous levels of protein.
<b>Purification</b>	COAA1 antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Collagen alpha-1 X chain
<b>Molecular Weight</b>	74 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid form in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml

<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:21850MIM:120110</a>
<b>Alternative Names</b>	Collagen alpha-1 X chain
<b>Function</b>	Type X collagen is a product of hypertrophic chondrocytes and has been localized to presumptive mineralization zones of hyaline cartilage.
<b>Cellular Localization</b>	Secreted, extracellular space, extracellular matrix
<b>Post-translational Modifications</b>	Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)