


## **Anti-ACE1** antibody



**Description** Rabbit polyclonal to ACE1.

Model STJ91441

**Host** Rabbit

**Reactivity** Human, Mouse, Rat

**Applications** ELISA, IHC, WB

**Immunogen** Synthesized peptide derived from human ACE1.

Immunogen Region Internal

**Gene ID** <u>1636</u>

Gene Symbol ACE

**Dilution range** WB 1:500-1:2000IHC 1:100-1:300ELISA 1:10000

**Specificity** ACE1 Polyclonal Antibody detects endogenous levels of ACE1 protein.

**Tissue Specificity** Ubiquitously expressed, with highest levels in lung, kidney, heart,

gastrointestinal system and prostate. Isoform Testis-specific is expressed in

spermatocytes and adult testis.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** Angiotensin-converting enzyme ACE Dipeptidyl carboxypeptidase I Kininase

II CD antigen CD143 Angiotensin-converting enzyme, soluble form

Molecular Weight 165 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:2707OMIM:106180

Alternative Names Angiotensin-converting enzyme ACE Dipeptidyl carboxypeptidase I Kininase

II CD antigen CD143 Angiotensin-converting enzyme, soluble form

**Function** Converts angiotensin I to angiotensin II by release of the terminal His-Leu,

this results in an increase of the vasoconstrictor activity of angiotensin. Also able to inactivate bradykinin, a potent vasodilator. Has also a glycosidase activity which releases GPI-anchored proteins from the membrane by cleaving

the mannose linkage in the GPI moiety.

**Cellular Localization** Angiotensin-converting enzyme, soluble form: Secreted.. Cell membrane.

Single-pass type I membrane protein. Cytoplasm. Detected in both cell

membrane and cytoplasm in neurons.

Post-translational

**Modifications** 

Phosphorylated by CK2 on Ser-1299; which allows membrane retention.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com