


Anti-Actin alpha antibody



Description	Rabbit polyclonal to Actin alpha3.
Model	STJ91463
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human Actin alpha3
Immunogen Region	1-80 aa, N-terminal
Gene ID	72
Gene Symbol	ACTG2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000
Specificity	Actin alpha3 Polyclonal Antibody detects endogenous levels of Actin alpha3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Actin, gamma-enteric smooth muscle Alpha-actin-3 Gamma-2-actin Smooth muscle gamma-actin
Molecular Weight	45 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:145OMIM:102545
Alternative Names	Actin, gamma-enteric smooth muscle Alpha-actin-3 Gamma-2-actin Smooth muscle gamma-actin
Function	Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.
Cellular Localization	Cytoplasm, cytoskeleton.
Post-translational Modifications	Oxidation of Met-45 and Met-48 by MICALs (MICAL1, MICAL2 or MICAL3) to form methionine sulfoxide promotes actin filament depolymerization. MICAL1 and MICAL2 produce the (R)-S-oxide form. The (R)-S-oxide form is reverted by MSRB1 and MSRB2, which promote actin repolymerization . Monomethylation at Lys-85 (K84me1) regulates actin-myosin interaction and actomyosin-dependent processes. Demethylation by ALKBH4 is required for maintaining actomyosin dynamics supporting normal cleavage furrow ingression during cytokinesis and cell migration . (Microbial infection) Monomeric actin is cross-linked by V.cholerae toxins RtxA and VgrG1 in case of infection: bacterial toxins mediate the cross-link between Lys-51 of one monomer and Glu-271 of another actin monomer, resulting in formation of highly toxic actin oligomers that cause cell rounding . The toxin can be highly efficient at very low concentrations by acting on formin homology family proteins: toxic actin oligomers bind with high affinity to formins and adversely affect both nucleation and elongation abilities of formins, causing their potent inhibition in both profilin-dependent and independent manners .