


Anti-ALS2CR8 antibody



Description Rabbit polyclonal to ALS2CR8.

Model STJ91571

Host Rabbit

Reactivity Human

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human ALS2CR8

Immunogen Region 280-360 aa, Internal

Gene ID <u>79800</u>

Gene Symbol <u>CARF</u>

Dilution range WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000

Specificity ALS2CR8 Polyclonal Antibody detects endogenous levels of ALS2CR8

protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Calcium-responsive transcription factor Amyotrophic lateral sclerosis 2

chromosomal region candidate gene 8 protein Calcium-response factor CaRF

Testis development protein NYD-SP24

Molecular Weight 85 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:14435OMIM:607586</u>

Alternative Names Calcium-responsive transcription factor Amyotrophic lateral sclerosis 2

chromosomal region candidate gene 8 protein Calcium-response factor CaRF

Testis development protein NYD-SP24

Function Acts as a transcriptional activator that mediates the calcium- and neuron-

selective induction of BDNF exon III transcription. Binds to the consensus

calcium-response element CaRE1 5'-CTATTTCGAG-3' sequence.

Sequence and Domain Family The N-terminus is necessary for DNA-binding. The C-terminus is necessary

for transcriptional activation.

Cellular Localization Nucleus

St John's Laboratory Ltd F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com