

Anti-ARA54 antibody



Description	Rabbit polyclonal to ARA54.
Model	STJ91666
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human ARA54
Immunogen Region	330-410 aa, C-terminal
Gene ID	9604
Gene Symbol	RNF14
Dilution range	WB 1:500-1:2000ELISA 1:40000
Specificity	ARA54 Polyclonal Antibody detects endogenous levels of ARA54 protein.
Tissue Specificity	Widely expressed.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	E3 ubiquitin-protein ligase RNF14 Androgen receptor-associated protein 54 HFB30 RING finger protein 14 Triad2 protein
Molecular Weight	50 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:10058OMIM:605675
Alternative Names	E3 ubiquitin-protein ligase RNF14 Androgen receptor-associated protein 54 HFB30 RING finger protein 14 Triad2 protein
Function	Might act as an E3 ubiquitin-protein ligase which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes and then transfers it to substrates, which could be nuclear proteins. Could play a role as a coactivator for androgen- and, to a lesser extent, progesterone-dependent transcription.
Sequence and Domain Family	The N-terminal destruction box (D-box) acts as a recognition signal for degradation via the ubiquitin-proteasome pathway. The RING-type zinc finger is essential for the interaction with UBE2E2.; Members of the RBR family are atypical E3 ligases. They interact with the E2 conjugating enzyme UBE2L3 and function like HECT-type E3 enzymes: they bind E2s via the first RING domain, but require an obligate trans-thiolation step during the ubiquitin transfer, requiring a conserved cysteine residue in the second RING domain.
Cellular Localization	Cytoplasm Nucleus
Post-translational Modifications	RING-type zinc finger-dependent and UBE2E2-dependent autoubiquitination.