


Anti-ARA70 antibody



Description	Rabbit polyclonal to ARA70.
Model	STJ91667
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human ARA70
Immunogen Region	320-400 aa, Internal
Gene ID	8031
Gene Symbol	NCOA4
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	ARA70 Polyclonal Antibody detects endogenous levels of ARA70 protein.
Tissue Specificity	Widely expressed. Also detected in adipose tissues and in different cell lines. Isoform Beta is only expressed in testis.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Nuclear receptor coactivator 4 NCoA-4 Androgen receptor coactivator 70 kDa protein 70 kDa AR-activator 70 kDa androgen receptor coactivator Androgen receptor-associated protein of 70 kDa Ret-activating protein ELE1
Molecular Weight	70 kDa

Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:7671OMIM:601984
Alternative Names	Nuclear receptor coactivator 4 NCoA-4 Androgen receptor coactivator 70 kDa protein 70 kDa AR-activator 70 kDa androgen receptor coactivator Androgen receptor-associated protein of 70 kDa Ret-activating protein ELE1
Function	Enhances the androgen receptor transcriptional activity in prostate cancer cells. Ligand-independent coactivator of the peroxisome proliferator-activated receptor (PPAR) gamma.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com