


## Anti-ATF-1 antibody



**Description** Rabbit polyclonal to ATF-1.

Model STJ91743

**Host** Rabbit

**Reactivity** Human, Mouse

**Applications** ELISA, IHC, WB

**Immunogen** Synthesized peptide derived from human ATF-1 around the non-

phosphorylation site of S63.

Immunogen Region 1-80 aa

**Gene ID** <u>466</u>

Gene Symbol <u>ATF1</u>

**Dilution range** WB 1:500-1:2000IHC 1:100-1:300ELISA 1:5000

**Specificity** ATF-1 Polyclonal Antibody detects endogenous levels of ATF-1 protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** Cyclic AMP-dependent transcription factor ATF-1 cAMP-dependent

transcription factor ATF-1 Activating transcription factor 1 Protein TREB36

Molecular Weight 29 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:783OMIM:123803</u>

Alternative Names Cyclic AMP-dependent transcription factor ATF-1 cAMP-dependent

transcription factor ATF-1 Activating transcription factor 1 Protein TREB36

**Function** This protein binds the cAMP response element (CRE) (consensus: 5'-

GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Binds to the Tax-responsive element (TRE) of HTLV-I. Mediates PKA-induced stimulation of CRE-reporter genes. Represses the expression of FTH1 and other antioxidant detoxification genes. Triggers cell proliferation

and transformation.

Cellular Localization Nucleus.

Post-translational Pho Modifications phos

Phosphorylated at Ser-198 by HIPK2 in response to genotoxic stress. This phosphorylation promotes transcription repression of FTH1 and other antioxidant detoxification genes. The CDK3-mediated phosphorylation at

Ser-63 promotes its transactivation and transcriptional activities. Phosphorylated at Ser-63 by RPS6KA4 and RPS6KA5 in response to

mitogenic or stress stimuli.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com