


## **Anti-ATP5G3 antibody**





Description	Rabbit polyclonal to ATP5G3.

Model STJ91770

**Host** Rabbit

**Reactivity** Human, Rat

**Applications** ELISA, IF, IHC

ImmunogenSynthesized peptide derived from human ATP5G3

Immunogen Region 10-90 aa, N-terminal

**Gene ID** <u>518</u>

Gene Symbol ATP5G3

**Dilution range** IHC 1:100-1:300IF 1:200-1:1000ELISA 1:40000

**Specificity** ATP5G3 Polyclonal Antibody detects endogenous levels of ATP5G3 protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

Protein Name ATP synthase F 0 complex subunit C3, mitochondrial ATP synthase lipid-

binding protein ATP synthase proteolipid P3 ATP synthase proton-

transporting mitochondrial F 0 complex subunit C3 ATPase protein 9 ATPase

subun

Molecular Weight 14.693 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:843OMIM:602736</u>

Alternative Names ATP synthase F 0 complex subunit C3, mitochondrial ATP synthase lipid-

binding protein ATP synthase proteolipid P3 ATP synthase proton-

transporting mitochondrial F 0 complex subunit C3 ATPase protein 9 ATPase

subun

**Function** Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex

V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain. A homomeric c-ring of

probably 10 subunits is part of the complex rotary element.

**Cellular Localization** Mitochondrion membrane. Multi-pass membrane protein.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com