


Anti-AW-1 antibody



Description	Rabbit polyclonal to AW-1.
Model	STJ91790
Host	Rabbit
Reactivity	Human
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human AW-1
Immunogen Region	180-260 aa, Internal
Gene ID	7727
Gene Symbol	ZNF174
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000
Specificity	AW-1 Polyclonal Antibody detects endogenous levels of AW-1 protein.
Tissue Specificity	Expressed in a variety of organs, but most strongly in adult testis and ovary followed by small intestine, colon, prostate, thymus, spleen, pancreas, skeletal muscle, heart, brain and kidney. Also expressed in umbilical vein endothelial cells, foreskin fibroblast and Hep-G2 cells.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Zinc finger protein 174 AW-1 Zinc finger and SCAN domain-containing protein 8

Molecular Weight	46 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:12963 OMIM:603900
Alternative Names	Zinc finger protein 174 AW-1 Zinc finger and SCAN domain-containing protein 8
Function	Transcriptional repressor.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com