


Anti-Bag-4 antibody



Description	Rabbit polyclonal to Bag-4.
Model	STJ91806
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human Bag-4.
Immunogen Region	C-terminal
Gene ID	9530
Gene Symbol	BAG4
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	Bag-4 Polyclonal Antibody detects endogenous levels of Bag-4 protein.
Tissue Specificity	Ubiquitous.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	BAG family molecular chaperone regulator 4 BAG-4 Bcl-2-associated athanogene 4 Silencer of death domains
Molecular Weight	49 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:940OMIM:603884
Alternative Names	BAG family molecular chaperone regulator 4 BAG-4 Bcl-2-associated athanogene 4 Silencer of death domains
Function	Inhibits the chaperone activity of HSP70/HSC70 by promoting substrate release . Prevents constitutive TNFRSF1A signaling. Negative regulator of PRKN translocation to damaged mitochondria.
Cellular Localization	Cytoplasm.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com