

Anti-C1q-C antibody



Description	Rabbit polyclonal to C1q-C.
Model	STJ91930
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human C1q-C
Immunogen Region	50-130 aa, Internal
Gene ID	714
Gene Symbol	C1QC
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	C1q-C Polyclonal Antibody detects endogenous levels of C1q-C protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Complement C1q subcomponent subunit C
Molecular Weight	30 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:1245OMIM:120575
Alternative Names	Complement C1q subcomponent subunit C
Function	C1q associates with the proenzymes C1r and C1s to yield C1, the first component of the serum complement system. The collagen-like regions of C1q interact with the Ca(2+)-dependent C1r(2)C1s(2) proenzyme complex, and efficient activation of C1 takes place on interaction of the globular heads of C1q with the Fc regions of IgG or IgM antibody present in immune complexes.
Cellular Localization	Secreted.
Post-translational Modifications	O-linked glycans consist of Glc-Gal disaccharides bound to the oxygen atom of post-translationally added hydroxyl groups.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com