

## Anti-Cadherin-22 antibody



**Description** Rabbit polyclonal to Cadherin-22.

Model STJ91963

**Host** Rabbit

**Reactivity** Human, Mouse, Rat

**Applications** ELISA, IF, IHC, WB

Immunogen Synthesized peptide derived from human Cadherin-22

**Immunogen Region** 80-160 aa, Internal

**Gene ID** <u>64405</u>

Gene Symbol CDH22

**Dilution range** WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000

Specificity Cadherin-22 Polyclonal Antibody detects endogenous levels of Cadherin-22

protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** Cadherin-22 Pituitary and brain cadherin PB-cadherin

Molecular Weight 84 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:13251OMIM:609920</u>

Alternative Names Cadherin-22 Pituitary and brain cadherin PB-cadherin

**Function** Cadherins are calcium-dependent cell adhesion proteins. They preferentially

interact with themselves in a homophilic manner in connecting cells;

cadherins may thus contribute to the sorting of heterogeneous cell types. PB-cadherins may have a role in the morphological organization of pituitary gland

and brain tissues.

Sequence and Domain Family Three calcium ions are usually bound at the interface of each cadherin domain

and rigidify the connections, imparting a strong curvature to the full-length

ectodomain.

Cellular Localization Cell membrane

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

W http://www.stjohnslabs.com/ E info@stjohnslabs.com

**T** +44 (0)208 223 3081 **E** info@stje