


Anti-CaMKI beta/gamma/delta antibody





Description Rabbit polyclonal to CaMKIIbeta/gamma/delta.

Model STJ91991

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human CaMKIIbeta/gamma/delta around

the non-phosphorylation site of T287.

Immunogen Region 230-310 aa

Gene ID <u>816</u>

Gene Symbol CAMK2B

Dilution range WB 1:500-1:2000IHC 1:100-1:300ELISA 1:5000

Specificity CaMKIIbeta/gamma/delta Polyclonal Antibody detects endogenous levels of

CaMKIIbeta/gamma/delta protein.

Tissue Specificity Widely expressed. Expressed in adult and fetal brain. Expression is slightly

lower in fetal brain. Expressed in skeletal muscle.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Calcium/calmodulin-dependent protein kinase type II subunit beta CaM

kinase II subunit beta CaMK-II subunit beta

Molecular Weight 50/65 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:1461OMIM:607707

Alternative Names Calcium/calmodulin-dependent protein kinase type II subunit beta CaM

kinase II subunit beta CaMK-II subunit beta

Function Calcium/calmodulin-dependent protein kinase that functions autonomously

after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in dendritic spine and synapse formation, neuronal plasticity and regulation of sarcoplasmic reticulum Ca(2+) transport in skeletal muscle. In neurons, plays an essential structural role in the reorganization of the actin cytoskeleton during plasticity by binding and bundling actin filaments in a kinase-

independent manner. This structural function is required for correct targeting of CaMK2A, which acts downstream of NMDAR to promote dendritic spine and synapse formation and maintain synaptic plasticity which enables long-term potentiation (LTP) and hippocampus-dependent learning. In developing hippocampal neurons, promotes arborization of the dendritic tree and in mature neurons, promotes dendritic remodeling. Participates in the modulation of skeletal muscle function in response to exercise. In slow-twitch muscles, is involved in regulation of sarcoplasmic reticulum (SR) Ca(2+) transport and in fast-twitch muscle participates in the control of Ca(2+) release from the SR through phosphorylation of triadin, a ryanodine receptor-coupling factor, and

Sequence and Domain Family The CAMK2 protein kinases contain a unique C-terminal subunit association

domain responsible for oligomerization.

Cellular Localization Cytoplasm, cytoskeleton Cytoplasm, cytoskeleton, microtubule organizing

center, centrosome Sarcoplasmic reticulum membrane. In slow-twitch muscle,

phospholamban (PLN/PLB), an endogenous inhibitor of SERCA2A/ATP2A2.

evenly distributed between longitudinal SR and junctional SR.

Post-translational Autophosphorylation of Thr-287 following activation by Ca(2+)/calmodulin.

Modifications Phosphorylation of Thr-287 locks the kinase into an activated state.