


## Anti-Cbl antibody



<b>Description</b>	Rabbit polyclonal to Cbl.
<b>Model</b>	STJ92055
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, IF, IHC, WB
<b>Immunogen</b>	Synthesized peptide derived from human Cbl around the non-phosphorylation site of Y774.
<b>Immunogen Region</b>	710-790 aa
<b>Gene ID</b>	<a href="#">867</a>
<b>Gene Symbol</b>	<a href="#">CBL</a>
<b>Dilution range</b>	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000
<b>Specificity</b>	Cbl Polyclonal Antibody detects endogenous levels of Cbl protein.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	E3 ubiquitin-protein ligase CBL Casitas B-lineage lymphoma proto-oncogene Proto-oncogene c-Cbl RING finger protein 55 RING-type E3 ubiquitin transferase CBL Signal transduction protein CBL
<b>Molecular Weight</b>	55120 kDa
<b>Clonality</b>	Polyclonal

<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:1541OMIM:165360</a>
<b>Alternative Names</b>	E3 ubiquitin-protein ligase CBL Casitas B-lineage lymphoma proto-oncogene Proto-oncogene c-Cbl RING finger protein 55 RING-type E3 ubiquitin transferase CBL Signal transduction protein CBL
<b>Function</b>	Adapter protein that functions as a negative regulator of many signaling pathways that are triggered by activation of cell surface receptors. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, EGFR, CSF1R, EPHA8 and KDR and terminates signaling. Recognizes membrane-bound HCK, SRC and other kinases of the SRC family and mediates their ubiquitination and degradation. Participates in signal transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast differentiation and apoptosis. Essential for osteoclastic bone resorption. The 'Tyr-731' phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function. May be functionally coupled with the E2 ubiquitin-protein ligase UB2D3.
<b>Sequence and Domain Family</b>	The RING-type zinc finger domain mediates binding to an E2 ubiquitin-conjugating enzyme.; The N-terminus is composed of the phosphotyrosine binding (PTB) domain, a short linker region and the RING-type zinc finger. The PTB domain, which is also called TKB (tyrosine kinase binding) domain, is composed of three different subdomains: a four-helix bundle (4H), a calcium-binding EF hand and a divergent SH2 domain.
<b>Cellular Localization</b>	Cytoplasm. Cell membrane. Colocalizes with FGFR2 in lipid rafts at the cell membrane.
<b>Post-translational Modifications</b>	Phosphorylated on tyrosine residues by ALK, EGFR, SYK, FYN and ZAP70 . Phosphorylated on tyrosine residues in response to FLT1 and KIT signaling. Phosphorylated on tyrosine residues by INSR and FGR. Phosphorylated on several tyrosine residues by constitutively activated FGFR3. Not phosphorylated at Tyr-731 by FGFR3. Phosphorylated on tyrosine residues by activated CSF1R, PDGFRA and PDGFRB. Phosphorylated on tyrosine residues by HCK. Ubiquitinated, leading to its degradation via the proteasome.