


Anti-CD40 antibody



Description	Rabbit polyclonal to CD40.
Model	STJ92128
Host	Rabbit
Reactivity	Human, Mouse, Simian
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human CD40
Immunogen Region	200-280 aa, C-terminal
Gene ID	958
Gene Symbol	CD40
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:10000
Specificity	CD40 Polyclonal Antibody detects endogenous levels of CD40 protein.
Tissue Specificity	B-cells and in primary carcinomas.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor superfamily member 5 B-cell surface antigen CD40 Bp50 CD40L receptor CDw40 CD antigen CD40
Molecular Weight	30 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11919OMIM:109535
Alternative Names	Tumor necrosis factor receptor superfamily member 5 B-cell surface antigen CD40 Bp50 CD40L receptor CDw40 CD antigen CD40
Function	Receptor for TNFSF5/CD40LG. Transduces TRAF6- and MAP3K8-mediated signals that activate ERK in macrophages and B cells, leading to induction of immunoglobulin secretion.
Cellular Localization	Isoform I: Cell membrane. Single-pass type I membrane protein.. Isoform II: Secreted.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com