


Anti-Cdc42EP2 antibody



Description Rabbit polyclonal to Cdc42EP2.

Model STJ92173

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human Cdc42EP2

Immunogen Region 10-90 aa, N-terminal

Gene ID <u>10435</u>

Gene Symbol CDC42EP2

Dilution range WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000

Specificity Cdc42EP2 Polyclonal Antibody detects endogenous levels of Cdc42EP2

protein.

Tissue Specificity Highly expressed in the heart. Weakly expressed in the pancreas and liver.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Cdc42 effector protein 2 Binder of Rho GTPases 1

Molecular Weight 28 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:16263OMIM:606132</u>

Alternative Names Cdc42 effector protein 2 Binder of Rho GTPases 1

Function Probably involved in the organization of the actin cytoskeleton. May act

downstream of CDC42 to induce actin filament assembly leading to cell shape changes. Induces pseudopodia formation in fibroblasts in a CDC42-dependent

manner.

Sequence and Domain Family The CRIB domain mediates interaction with CDC42.

Cellular Localization Endomembrane system Cytoplasm, cytoskeleton

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/

E info@stjohnslabs.com