


Anti-CdcA3 antibody



Description	Rabbit polyclonal to CdcA3.
Model	STJ92184
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human CdcA3
Immunogen Region	190-270 aa, C-terminal
Gene ID	83461
Gene Symbol	CDCA3
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000
Specificity	CdcA3 Polyclonal Antibody detects endogenous levels of CdcA3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Cell division cycle-associated protein 3 Gene-rich cluster protein C8 Trigger of mitotic entry protein 1 TOME-1
Molecular Weight	28 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:14624OMIM:607749
Alternative Names	Cell division cycle-associated protein 3 Gene-rich cluster protein C8 Trigger of mitotic entry protein 1 TOME-1
Function	F-box-like protein which is required for entry into mitosis. Acts by participating in E3 ligase complexes that mediate the ubiquitination and degradation of WEE1 kinase at G2/M phase .
Sequence and Domain Family	The KEN box is required for the association with the APC/C-Cdh1 complex.
Cellular Localization	Cytoplasm, cytosol
Post-translational Modifications	Ubiquitinated and degraded by the APC/C-Cdh1 complex.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com