

## Anti-CIDE-3 antibody



<b>Description</b>	Rabbit polyclonal to CIDE-3.
--------------------	------------------------------

<b>Model</b>	STJ92292
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA, IF, IHC
<b>Immunogen</b>	Synthesized peptide derived from human CIDE-3
<b>Immunogen Region</b>	160-240 aa, C-terminal
<b>Gene ID</b>	<a href="#">63924</a>
<b>Gene Symbol</b>	<a href="#">CIDEc</a>
<b>Dilution range</b>	IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000
<b>Specificity</b>	CIDE-3 Polyclonal Antibody detects endogenous levels of CIDE-3 protein.
<b>Tissue Specificity</b>	Expressed mainly in adipose tissue, small intestine, heart, colon and stomach and, at lower levels, in brain, kidney and liver.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Cell death activator CIDE-3 Cell death-inducing DFFA-like effector protein C Fat-specific protein FSP27 homolog
<b>Molecular Weight</b>	26.754 kDa
<b>Clonality</b>	Polyclonal

<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:24229</a> <a href="#">OMIM:612120</a>
<b>Alternative Names</b>	Cell death activator CIDE-3 Cell death-inducing DFFA-like effector protein C Fat-specific protein FSP27 homolog
<b>Function</b>	Binds to lipid droplets and regulates their enlargement, thereby restricting lipolysis and favoring storage. At focal contact sites between lipid droplets, promotes directional net neutral lipid transfer from the smaller to larger lipid droplets. The transfer direction may be driven by the internal pressure difference between the contacting lipid droplet pair. Its role in neutral lipid transfer and lipid droplet enlargement is activated by the interaction with PLIN1. May act as a CEBPB coactivator in the white adipose tissue to control the expression of a subset of CEBPB downstream target genes, including SOCS1, SOCS3, TGFB1, TGFBR1, ID2 and XDH. When overexpressed in preadipocytes, induces apoptosis or increases cell susceptibility to apoptosis induced by serum deprivation or TGFB treatment. As mature adipocytes, that express high CIDEC levels, are quite resistant to apoptotic stimuli, the physiological significance of its role in apoptosis is unclear. May play a role in the modulation of the response to osmotic stress by preventing NFAT5 to translocate into the nucleus and activate its target genes expression.
<b>Sequence and Domain Family</b>	The CIDE-N domain is involved in homodimerization which is crucial for its function in promoting lipid exchange and transfer.
<b>Cellular Localization</b>	Nucleus Endoplasmic reticulum. Lipid droplet. Diffuses quickly on lipid droplet surface, but becomes trapped and clustered at lipid droplet contact sites, thereby enabling its rapid enrichment at lipid droplet contact sites.
<b>Post-translational Modifications</b>	Ubiquitinated and targeted to proteasomal degradation, resulting in a short half-life. Protein stability depends on triacylglycerol synthesis, fatty acid availability and lipid droplet formation .