

Anti-c-Kit antibody



Description	Rabbit polyclonal to c-Kit.
--------------------	-----------------------------

Model	STJ92298
Host	Rabbit
Reactivity	Human, Mouse, Rat, Simian
Applications	ELISA, IF, WB
Immunogen	Synthesized peptide derived from human c-Kit around the non-phosphorylation site of Y721.
Immunogen Region	660-740 aa
Gene ID	3815
Gene Symbol	KIT
Dilution range	WB 1:500-1:2000IF 1:200-1:1000ELISA 1:10000
Specificity	c-Kit Polyclonal Antibody detects endogenous levels of c-Kit protein.
Tissue Specificity	Isoform 1 and isoform 2 are detected in spermatogonia and Leydig cells. Isoform 3 is detected in round spermatids, elongating spermatids and spermatozoa (at protein level). Widely expressed. Detected in the hematopoietic system, the gastrointestinal system, in melanocytes and in germ cells.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Mast/stem cell growth factor receptor Kit SCFR Piebald trait protein PBT

	Proto-oncogene c-Kit Tyrosine-protein kinase Kit p145 c-kit v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog CD antigen C
Molecular Weight	120 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:6342 OMIM:164920
Alternative Names	Mast/stem cell growth factor receptor Kit SCFR Piebald trait protein PBT Proto-oncogene c-Kit Tyrosine-protein kinase Kit p145 c-kit v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog CD antigen C
Function	<p>Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine KITLG/SCF and plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. In response to KITLG/SCF binding, KIT can activate several signaling pathways. Phosphorylates PIK3R1, PLCG1, SH2B2/APS and CBL. Activates the AKT1 signaling pathway by phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Activated KIT also transmits signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family members STAT1, STAT3, STAT5A and STAT5B. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KIT signaling is modulated by protein phosphatases, and by rapid internalization and degradation of the receptor. Activated KIT promotes phosphorylation of the protein phosphatases PTPN6/SHP-1 and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B. Promotes phosphorylation of PIK3R1, CBL, CRK (isoform Crk-II), LYN, MAPK1/ERK2 and/or MAPK3/ERK1, PLCG1, SRC and SHC1.</p>
Cellular Localization	<p>Isoform 1: Cell membrane. Single-pass type I membrane protein.. Isoform 2: Cell membrane. Single-pass type I membrane protein.. Isoform 3: Cytoplasm. Detected in the cytoplasm of spermatozoa, especially in the equatorial and subacrosomal region of the sperm head.</p>
Post-translational Modifications	<p>Ubiquitinated by SOCS6. KIT is rapidly ubiquitinated after autophosphorylation induced by KITLG/SCF binding, leading to internalization and degradation. Autophosphorylated on tyrosine residues. KITLG/SCF binding enhances autophosphorylation. Isoform 1 shows low levels of tyrosine phosphorylation in the absence of added KITLG/SCF (in vitro). Kinase activity is down-regulated by phosphorylation on serine residues by protein kinase C family members. Phosphorylation at Tyr-568 is required for interaction with PTPN11/SHP-2, CRK (isoform Crk-II) and members of the SRC tyrosine-protein kinase family. Phosphorylation at Tyr-570 is required for interaction with PTPN6/SHP-1. Phosphorylation at Tyr-703, Tyr-823 and Tyr-936 is important for interaction with GRB2.</p>

Phosphorylation at Tyr-721 is important for interaction with PIK3R1.
Phosphorylation at Tyr-823 and Tyr-936 is important for interaction with GRB7.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com