


Anti-CstF-64 antibody



Description	Rabbit polyclonal to CstF-64.
Model	STJ92510
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human CstF-64
Immunogen Region	30-110 aa, N-terminal
Gene ID	1478
Gene Symbol	CSTF2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000
Specificity	CstF-64 Polyclonal Antibody detects endogenous levels of CstF-64 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Cleavage stimulation factor subunit 2 CF-1 64 kDa subunit Cleavage stimulation factor 64 kDa subunit CSTF 64 kDa subunit CstF-64
Molecular Weight	60 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:2484OMIM:300907
Alternative Names	Cleavage stimulation factor subunit 2 CF-1 64 kDa subunit Cleavage stimulation factor 64 kDa subunit CSTF 64 kDa subunit CstF-64
Function	One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs. This subunit is directly involved in the binding to pre-mRNAs .
Cellular Localization	Nucleus. Localized with DDX1 in cleavage bodies.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com