


Anti-C-TAK1 antibody



Description	Rabbit polyclonal to C-TAK1.
Model	STJ92512
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human C-TAK1
Immunogen Region	1-80 aa, N-terminal
Gene ID	4140
Gene Symbol	MARK3
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:40000
Specificity	C-TAK1 Polyclonal Antibody detects endogenous levels of C-TAK1 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	MAP/microtubule affinity-regulating kinase 3 C-TAK1 cTAK1 Cdc25C-associated protein kinase 1 ELKL motif kinase 2 EMK-2 Protein kinase STK10 Ser/Thr protein kinase PAR-1 Par-1a Serine/threonine-protein
Molecular Weight	85 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:6897OMIM:602678
Alternative Names	MAP/microtubule affinity-regulating kinase 3 C-TAK1 cTAK1 Cdc25C-associated protein kinase 1 ELKL motif kinase 2 EMK-2 Protein kinase STK10 Ser/Thr protein kinase PAR-1 Par-1a Serine/threonine-protein

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com