


Anti-Cyclin A1 antibody



Description	Rabbit polyclonal to Cyclin A1.
Model	STJ92533
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human Cyclin A1
Immunogen Region	380-460 aa, C-terminal
Gene ID	8900
Gene Symbol	CCNA1
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	Cyclin A1 Polyclonal Antibody detects endogenous levels of Cyclin A1 protein.
Tissue Specificity	Very high levels in testis and very low levels in brain. Also found in myeloid leukemia cell lines.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Cyclin-A1
Molecular Weight	52 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:1577OMIM:604036
Alternative Names	Cyclin-A1
Function	May be involved in the control of the cell cycle at the G1/S (start) and G2/M (mitosis) transitions. May primarily function in the control of the germline meiotic cell cycle and additionally in the control of mitotic cell cycle in some somatic cells.
Cellular Localization	Nucleus
Post-translational Modifications	Polyubiquitinated via 'Lys-11'-linked ubiquitin by the anaphase-promoting complex (APC/C), leading to its degradation by the proteasome. Deubiquitinated and stabilized by USP37 enables entry into S phase.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com