

Anti-Cyclin E2 antibody



Description	Rabbit polyclonal to Cyclin E2.
Model	STJ92544
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human Cyclin E2 around the non-phosphorylation site of T392.
Immunogen Region	330-410 aa
Gene ID	9134
Gene Symbol	CCNE2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000
Specificity	Cyclin E2 Polyclonal Antibody detects endogenous levels of Cyclin E2 protein.
Tissue Specificity	According to PubMed:9858585, highest levels of expression in adult testis, thymus and brain. Lower levels in placenta, spleen and colon. Consistently elevated levels in tumor-derived cells compared to non-transformed proliferating cells. According to PubMed:9840927: low levels in thymus, prostate, brain, skeletal muscle, and kidney. Elevated levels in lung. According to PubMed:9840943 highly expressed in testis, placenta, thymus and brain. In a lesser extent in small intestine and colon.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Note	For Research Use Only (RUO).
Protein Name	G1/S-specific cyclin-E2
Molecular Weight	46 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:15900MIM:603775
Alternative Names	G1/S-specific cyclin-E2
Function	Essential for the control of the cell cycle at the late G1 and early S phase.
Cellular Localization	Nucleus
Post-translational Modifications	Phosphorylation by CDK2 triggers its release from CDK2 and degradation via the ubiquitin proteasome pathway.