

## **Anti-CYP4V2** antibody



**Description** Rabbit polyclonal to CYP4V2.

Model STJ92601

**Host** Rabbit

**Reactivity** Human

**Applications** ELISA, IHC, WB

**Immunogen** Synthesized peptide derived from human CYP4V2.

Immunogen Region Internal

**Gene ID** <u>285440</u>

Gene Symbol <u>CYP4V2</u>

**Dilution range** WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000

**Specificity** CYP4V2 Polyclonal Antibody detects endogenous levels of CYP4V2 protein.

**Tissue Specificity** Broadly expressed. Detected in heart, brain, placenta, lung, liver, skeletal

muscle, kidney, pancreas, retina, retinal pigment epithelium (RPE) and

lymphocytes.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

Protein Name Cytochrome P450 4V2 Docosahexaenoic acid omega-hydroxylase CYP4V2

Molecular Weight 60 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:23198OMIM:210370</u>

Alternative Names Cytochrome P450 4V2 Docosahexaenoic acid omega-hydroxylase CYP4V2

**Function** Omega-hydroxylase that oxidizes medium-chain saturated fatty acids and

polyunsaturated omega-3 fatty acids, and which plays a role in fatty acid and

steroid metabolism in the eye . Catalyzes the omega-hydroxylation of

medium-chain saturated fatty acids such as laurate, myristate and palmitate in an NADPH-dependent pathway. The substrate specificity is higher for

myristate > laurate > palmitate (C14>C16>C12). Acts as a polyunsaturated omega-3 fatty acids hydroxylase by mediating oxidation of docosahexaenoate

(DHA) to 22-hydroxydocosahexaenoate . Also produces some 21-

hydroxydocosahexaenoate. Also converts eicosapentaenoate (EPA) to 20-

hydroxyeicosapentaenoate (20-OH-EPA).

Cellular Localization Endoplasmic reticulum membrane

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com