Anti-DPF2 antibody



Description	Rabbit polyclonal to DPF2.

Model	STJ92773
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human DPF2
Immunogen Region	120-200 aa, Internal
Gene ID	5977
Gene Symbol	DPF2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000
Specificity	DPF2 Polyclonal Antibody detects endogenous levels of DPF2 protein.
Tissue Specificity	Ubiquitous.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Zinc finger protein ubi-d4 Apoptosis response zinc finger protein BRG1-associated factor 45D BAF45D D4, zinc and double PHD fingers family 2 Protein requiem
Molecular Weight	44 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9964OMIM:601671
Alternative Names	Zinc finger protein ubi-d4 Apoptosis response zinc finger protein BRG1-associated factor 45D BAF45D D4, zinc and double PHD fingers family 2 Protein requiem
Function	May be a transcription factor required for the apoptosis response following survival factor withdrawal from myeloid cells. Might also have a role in the development and maturation of lymphoid cells.
Cellular Localization	Nucleus. Cytoplasm. 30% nuclear. 70% cytoplasmic.