


Anti-ELOVL3 antibody





Description Rabbit polyclonal to ELOVL3.

Model STJ92902

Host Rabbit

Reactivity Human

ELISA, IHC **Applications**

Synthesized peptide derived from human ELOVL3 **Immunogen**

1-80 aa, Internal **Immunogen Region**

Gene ID 83401

Gene Symbol ELOVL3

Dilution range IHC 1:100-1:300ELISA 1:20000

ELOVL3 Polyclonal Antibody detects endogenous levels of ELOVL3 protein. **Specificity**

Testis. **Tissue Specificity**

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Elongation of very long chain fatty acids protein 3 3-keto acyl-CoA synthase

> ELOVL3 Cold-inducible glycoprotein of 30 kDa ELOVL fatty acid elongase 3 ELOVL FA elongase 3 Very long chain 3-ketoacyl-CoA synthase 3 Very lo

Molecular Weight 31.5 kDa

Clonality Polyclonal **Conjugation** Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:18047OMIM:611815</u>

Alternative Names Elongation of very long chain fatty acids protein 3 3-keto acyl-CoA synthase

ELOVL3 Cold-inducible glycoprotein of 30 kDa ELOVL fatty acid elongase 3 ELOVL FA elongase 3 Very long chain 3-ketoacyl-CoA synthase 3 Very lo

Function Catalyzes the first and rate-limiting reaction of the four that constitute the

long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process, allows the addition of 2 carbons to the chain of long- and very long-chain fatty acids/VLCFAs per cycle. Condensing enzyme with higher activity toward C18 acyl-CoAs, especially C18:0 acyl-CoAs. May participate in the production of saturated and monounsaturated VLCFAs of different chain lengths that are involved in multiple biological processes as

precursors of membrane lipids and lipid mediators.

Sequence and Domain Family The C-terminal di-lysine motif may confer endoplasmic reticulum

localization.

Cellular Localization Endoplasmic reticulum membrane

Post-translational N-Glycosylated.

Modifications

St John's Laboratory Ltd F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com