


Anti-ERK 8 antibody



Description Rabbit polyclonal to ERK 8.

Model STJ92995

Host Rabbit

Reactivity Human, Mouse

Applications ELISA, IF, IHC, WB

Immunogen Synthesized peptide derived from human ERK 8 around the non-

phosphorylation site of T175/Y177.

Immunogen Region 110-190 aa

Gene ID 225689

Gene Symbol MAPK15

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000

Specificity ERK 8 Polyclonal Antibody detects endogenous levels of ERK 8 protein.

Tissue Specificity Widely expressed with a maximal expression in lung and kidney.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Mitogen-activated protein kinase 15 MAP kinase 15 MAPK 15 Extracellular

signal-regulated kinase 7 ERK-7 Extracellular signal-regulated kinase 8

ERK-8

Molecular Weight 60 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:24667</u>OMIM:NA

Alternative Names Mitogen-activated protein kinase 15 MAP kinase 15 MAPK 15 Extracellular

signal-regulated kinase 7 ERK-7 Extracellular signal-regulated kinase 8

ERK-8

Function In vitro, phosphorylates MBP.

Sequence and Domain Family The N-terminal region (1-20) is the minimal region necessary for

ubiquitination and further proteasomal degradation. The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP

kinases.

Post-translational Dually phosphorylated on Thr-175 and Tyr-177, which activates the enzyme.

Autophosphorylated on threonine and tyrosine residues in vitro.

Ubiquitinated. Ubiquitination may allow its tight kinase activity regulation

and rapid turnover. May be ubiquitinated by a SCF E3 ligase.

St John's Laboratory Ltd

Modifications

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com