

Anti-FDC-SP antibody



Description	Rabbit polyclonal to FDC-SP.
Model	STJ93052
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC
Immunogen	Synthesized peptide derived from human FDC-SP.
Immunogen Region	C-terminal
Gene ID	260436
Gene Symbol	FDCSP
Dilution range	IHC 1:100-1:300ELISA 1:40000
Specificity	FDC-SP Polyclonal Antibody detects endogenous levels of FDC-SP protein.
Tissue Specificity	Abundantly expressed in tonsil, lymph node, and trachea; strong expression in prostate; lower expression in thyroid, stomach, and colon.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Follicular dendritic cell secreted peptide FDC secreted protein FDC-SP
Molecular Weight	9.7 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:19215OMIM:607241
Alternative Names	Follicular dendritic cell secreted peptide FDC secreted protein FDC-SP
Function	Can bind to the surface of B-lymphoma cells, but not T-lymphoma cells, consistent with a function as a secreted mediator acting upon B-cells.
Cellular Localization	Secreted.
Post-translational Modifications	O-glycosylated with core 1 or possibly core 8 glycans.

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com