

Anti-FKRP antibody



Description Rabbit polyclonal to FKRP.

Model STJ93080

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, IF, WB

ImmunogenSynthesized peptide derived from human FKRP

Immunogen Region 10-90 aa, N-terminal

Gene ID <u>79147</u>

Gene Symbol <u>FKRP</u>

Dilution range WB 1:500-1:2000IF 1:200-1:1000ELISA 1:20000

Specificity FKRP Polyclonal Antibody detects endogenous levels of FKRP protein.

Tissue Specificity Expressed predominantly in skeletal muscle, placenta, and heart and relatively

weakly in brain, lung, liver kidney and pancreas.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Fukutin-related protein

Molecular Weight 50 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:17997OMIM:606596</u>

Alternative Names Fukutin-related protein

Function Transferase involved in the biosynthesis of the phosphorylated O-mannosyl

trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-

dystroglycan (DAG1), which is required for binding laminin G-like domain-

containing extracellular proteins with high affinity.

Cellular Localization Golgi apparatus membrane Rough endoplasmic reticulum. According to some

studies the N-terminal hydrophobic domain is cleaved after translocation to the Golgi apparatus and the protein is secreted. Localization at the cell membrane may require the presence of dystroglycan. At the Golgi apparatus

localizes to the middle-to-trans-cisternae, as assessed by MG160

colocalization. Detected in rough endoplasmic reticulum in myocytes. In general, mutants associated with severe clinical phenotypes are retained

within the endoplasmic reticulum.

Post-translational

Modifications

N-glycosylated.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com