


Anti-Frizzled-1 antibody



Description	Rabbit polyclonal to Frizzled-1.
Model	STJ93138
Host	Rabbit
Reactivity	Human
Applications	ELISA, IF, WB
Immunogen	Synthesized peptide derived from human Frizzled-1
Immunogen Region	30-110 aa, N-terminal
Gene ID	8321
Gene Symbol	FZD1
Dilution range	WB 1:500-1:2000IF 1:200-1:1000ELISA 1:5000
Specificity	Frizzled-1 Polyclonal Antibody detects endogenous levels of Frizzled-1 protein.
Tissue Specificity	Expressed in adult heart, placenta, lung, kidney, pancreas, prostate, and ovary and in fetal lung and kidney.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Frizzled-1 Fz-1 hFz1 FzE1
Molecular Weight	71 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4038OMIM:603408
Alternative Names	Frizzled-1 Fz-1 hFz1 FzE1
Function	Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Activated by Wnt3A, Wnt3, Wnt1 and to a lesser extent Wnt2, but not by Wnt4, Wnt5A, Wnt5B, Wnt6, Wnt7A or Wnt7B.
Sequence and Domain Family	Lys-Thr-X-X-X-Trp motif interacts with the PDZ domain of Dvl (Disheveled) family members and is involved in the activation of the Wnt/beta-catenin signaling pathway. The FZ domain is involved in binding with Wnt ligands.
Cellular Localization	Membrane. Multi-pass membrane protein. Cell membrane
Post-translational Modifications	Ubiquitinated by ZNRF3, leading to its degradation by the proteasome.