

## Anti-Granzyme A antibody



**Description** Rabbit polyclonal to Granzyme A.

Model STJ93413

**Host** Rabbit

**Reactivity** Human

**Applications** ELISA, IF, IHC

Immunogen Synthesized peptide derived from human Granzyme A

**Immunogen Region** 30-110 aa, Internal

**Gene ID** <u>3001</u>

Gene Symbol GZMA

**Dilution range** IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000

**Specificity** Granzyme A Polyclonal Antibody detects endogenous levels of Granzyme A

protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

**Protein Name** Granzyme A CTL tryptase Cytotoxic T-lymphocyte proteinase 1

Fragmentin-1 Granzyme-1 Hanukkah factor H factor HF

Molecular Weight 28.969 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

1 mg/ml Concentration

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction** 

**Database Links** HGNC:4708OMIM:140050

**Alternative Names** Granzyme A CTL tryptase Cytotoxic T-lymphocyte proteinase 1

Fragmentin-1 Granzyme-1 Hanukkah factor H factor HF

**Function** Abundant protease in the cytosolic granules of cytotoxic T-cells and NK-cells

> which activates caspase-independent cell death with morphological features of apoptosis when delivered into the target cell through the immunological synapse. It cleaves after Lys or Arg. Cleaves APEX1 after 'Lys-31' and destroys its oxidative repair activity. Cleaves the nucleosome assembly protein SET after 'Lys-189', which disrupts its nucleosome assembly activity and allows the SET complex to translocate into the nucleus to nick and

degrade the DNA.

**Cellular Localization** Isoform alpha: Secreted. Cytoplasmic granule.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

W http://www.stjohnslabs.com/ T+44 (0)208 223 3081 E info@stjohnslabs.com