


## Anti-Hrs antibody



<b>Description</b>	Rabbit polyclonal to Hrs.
<b>Model</b>	STJ93593
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, IF, IHC, WB
<b>Immunogen</b>	Synthesized peptide derived from human Hrs around the non-phosphorylation site of Y334.
<b>Immunogen Region</b>	270-350 aa
<b>Gene ID</b>	<a href="#">9146</a>
<b>Gene Symbol</b>	<a href="#">HGS</a>
<b>Dilution range</b>	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000
<b>Specificity</b>	Hrs Polyclonal Antibody detects endogenous levels of Hrs protein.
<b>Tissue Specificity</b>	Ubiquitous expression in adult and fetal tissues with higher expression in testis and peripheral blood leukocytes.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Hepatocyte growth factor-regulated tyrosine kinase substrate Hrs Protein pp110
<b>Molecular Weight</b>	80 kDa

<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:48970MIM:604375</a>
<b>Alternative Names</b>	Hepatocyte growth factor-regulated tyrosine kinase substrate Hrs Protein pp110
<b>Function</b>	Involved in intracellular signal transduction mediated by cytokines and growth factors. When associated with STAM, it suppresses DNA signaling upon stimulation by IL-2 and GM-CSF. Could be a direct effector of PI3-kinase in vesicular pathway via early endosomes and may regulate trafficking to early and late endosomes by recruiting clathrin. May concentrate ubiquitinated receptors within clathrin-coated regions. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with STAM (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes. May contribute to the efficient recruitment of SMADs to the activin receptor complex. Involved in receptor recycling via its association with the CART complex, a multiprotein complex required for efficient transferrin receptor recycling but not for EGFR degradation.
<b>Sequence and Domain Family</b>	Has a double-sided UIM that can bind 2 ubiquitin molecules, one on each side of the helix.; The FYVE-type zinc finger domain mediates interactions with phosphatidylinositol 3-phosphate in membranes of early endosomes and penetrates bilayers. The FYVE domain insertion into PtdIns(3)P-enriched membranes is substantially increased in acidic conditions.
<b>Cellular Localization</b>	Cytoplasm Early endosome membrane Endosome, multivesicular body membrane. Colocalizes with UBQLN1 in ubiquitin-rich cytoplasmic aggregates that are not endocytic compartments.
<b>Post-translational Modifications</b>	Phosphorylated on Tyr-334. A minor site of phosphorylation on Tyr-329 is detected . Phosphorylation occurs in response to EGF, IL-2, GM-CSF and HGF. Ubiquitinated by ITCH.