


Anti-ICAD antibody



Description	Rabbit polyclonal to ICAD.
Model	STJ93633
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human ICAD
Immunogen Region	120-200 aa, Internal
Gene ID	1676
Gene Symbol	DFFA
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000
Specificity	ICAD Polyclonal Antibody detects endogenous levels of ICAD protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	DNA fragmentation factor subunit alpha DNA fragmentation factor 45 kDa subunit DFF-45 Inhibitor of CAD ICAD
Molecular Weight	36 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:2772OMIM:601882
Alternative Names	DNA fragmentation factor subunit alpha DNA fragmentation factor 45 kDa subunit DFF-45 Inhibitor of CAD ICAD
Function	Inhibitor of the caspase-activated DNase (DFF40).
Cellular Localization	Cytoplasm.
Post-translational Modifications	Caspase-3 cleaves DFF45 at 2 sites to generate an active factor.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com