

Anti-IL-12B antibody



Description	Rabbit polyclonal to IL-12B.
Model	STJ93677
Host	Rabbit
Reactivity	Human
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human IL-12B.
Immunogen Region	Internal
Gene ID	3593
Gene Symbol	IL12B
Dilution range	WB 1:500-1:2000ELISA 1:5000
Specificity	IL-12B Polyclonal Antibody detects endogenous levels of IL-12B protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Interleukin-12 subunit beta IL-12B Cytotoxic lymphocyte maturation factor 40 kDa subunit CLMF p40 IL-12 subunit p40 NK cell stimulatory factor chain 2 NKSF2
Molecular Weight	40 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:59700MIM:161561
Alternative Names	Interleukin-12 subunit beta IL-12B Cytotoxic lymphocyte maturation factor 40 kDa subunit CLMF p40 IL-12 subunit p40 NK cell stimulatory factor chain 2 NKSF2
Function	Cytokine that can act as a growth factor for activated T and NK cells, enhance the lytic activity of NK/lymphokine-activated killer cells, and stimulate the production of IFN-gamma by resting PBMC. Associates with IL23A to form the IL-23 interleukin, a heterodimeric cytokine which functions in innate and adaptive immunity. IL-23 may constitute with IL-17 an acute response to infection in peripheral tissues. IL-23 binds to a heterodimeric receptor complex composed of IL12RB1 and IL23R, activates the Jak-Stat signaling cascade, stimulates memory rather than naive T-cells and promotes production of proinflammatory cytokines. IL-23 induces autoimmune inflammation and thus may be responsible for autoimmune inflammatory diseases and may be important for tumorigenesis.
Cellular Localization	Secreted.
Post-translational Modifications	Known to be C-mannosylated in the recombinant protein; it is not yet known for sure if the wild-type protein is also modified.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com