

## Anti-KKIAMRE antibody

---



<b>Description</b>	Rabbit polyclonal to KKIAMRE.
<b>Model</b>	STJ93847
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA, IHC, WB
<b>Immunogen</b>	Synthesized peptide derived from human KKIAMRE
<b>Immunogen Region</b>	180-260 aa, Internal
<b>Gene ID</b>	<a href="#">8999</a>
<b>Gene Symbol</b>	<a href="#">CDKL2</a>
<b>Dilution range</b>	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:10000
<b>Specificity</b>	KKIAMRE Polyclonal Antibody detects endogenous levels of KKIAMRE protein.
<b>Tissue Specificity</b>	Expressed in testis and kidney, and at lower level in brain and lung.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Cyclin-dependent kinase-like 2 Protein kinase p56 KKIAMRE Serine/threonine-protein kinase KKIAMRE
<b>Molecular Weight</b>	56 kDa
<b>Clonality</b>	Polyclonal

<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:1782OMIM:603442</a>
<b>Alternative Names</b>	Cyclin-dependent kinase-like 2 Protein kinase p56 KKIAMRE Serine/threonine-protein kinase KKIAMRE
<b>Sequence and Domain Family</b>	The [NKR]KIAxRE motif seems to be a cyclin-binding region.
<b>Cellular Localization</b>	Cytoplasm Nucleus

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580  
**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>  
**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)