


Anti-NDRG4 antibody



| | |
|--------------------|-----------------------------|
| Description | Rabbit polyclonal to NDRG4. |
|--------------------|-----------------------------|

| | |
|---------------------------|--|
| Model | STJ94371 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Applications | ELISA, IHC, WB |
| Immunogen | Synthesized peptide derived from human NDRG4 |
| Immunogen Region | 160-240 aa, Internal |
| Gene ID | 65009 |
| Gene Symbol | NDRG4 |
| Dilution range | WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000 |
| Specificity | NDRG4 Polyclonal Antibody detects endogenous levels of NDRG4 protein. |
| Tissue Specificity | Expressed predominantly in brain and heart (at protein level). In the brain, detected in astrocytes. Isoform 1 and isoform 2 are only expressed in brain. Isoform 3 is expressed in both heart and brain. Up-regulated in glioblastoma multiforme cells. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Note | For Research Use Only (RUO). |
| Protein Name | Protein NDRG4 Brain development-related molecule 1 N-myc downstream-regulated gene 4 protein Vascular smooth muscle cell-associated protein 8 SMAP-8 |

| | |
|---|---|
| Molecular Weight | 34 kDa |
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Isotype | IgG |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Concentration | 1 mg/ml |
| Storage Instruction | Store at -20°C, and avoid repeat freeze-thaw cycles. |
| Database Links | HGNC:14466 OMIM:614463 |
| Alternative Names | Protein NDRG4 Brain development-related molecule 1 N-myc downstream-regulated gene 4 protein Vascular smooth muscle cell-associated protein 8 SMAP-8 |
| Function | Contributes to the maintenance of intracerebral BDNF levels within the normal range, which is necessary for the preservation of spatial learning and the resistance to neuronal cell death caused by ischemic stress. May enhance growth factor-induced ERK1 and ERK2 phosphorylation, including that induced by PDGF and FGF. May attenuate NGF-promoted ELK1 phosphorylation in a microtubule-dependent manner. |
| Cellular Localization | Cytoplasm, cytosol |
| Post-translational Modifications | Phosphorylated in an aortic smooth muscle cell line, following PDGF treatment. |

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com