

Anti-PBF antibody



Description Rabbit polyclonal to PBF.

Model STJ94974

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, IF, IHC, WB

ImmunogenSynthesized peptide derived from human PBF

Immunogen Region 90-170 aa, Internal

Gene ID <u>754</u>

Gene Symbol PTTG1IP

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000

Specificity PBF Polyclonal Antibody detects endogenous levels of PBF protein.

Tissue Specificity Ubiquitous.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Pituitary tumor-transforming gene 1 protein-interacting protein Pituitary

tumor-transforming gene protein-binding factor PBF PTTG-binding factor

Molecular Weight 20 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:13524OMIM:603784</u>

Alternative Names Pituitary tumor-transforming gene 1 protein-interacting protein Pituitary

tumor-transforming gene protein-binding factor PBF PTTG-binding factor

Function May facilitate PTTG1 nuclear translocation.

Cellular Localization Membrane Cytoplasm Nucleus. According to PubMed:10781616, it is found

in the cytoplasm and the nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/

E info@stjohnslabs.com