


Anti-PKA alpha reg antibody



Description	Rabbit polyclonal to PKA Ialpha reg.
Model	STJ95111
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human PKA Ialpha reg
Immunogen Region	240-320 aa, C-terminal
Gene ID	5573
Gene Symbol	PRKAR1A
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000
Specificity	PKA Ialpha reg Polyclonal Antibody detects endogenous levels of PKA Ialpha reg protein.
Tissue Specificity	Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	cAMP-dependent protein kinase type I-alpha regulatory subunit Tissue-specific extinguisher 1 TSE1

Molecular Weight	43 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9388OMIM:101800
Alternative Names	cAMP-dependent protein kinase type I-alpha regulatory subunit Tissue-specific extinguisher 1 TSE1
Function	Regulatory subunit of the cAMP-dependent protein kinases involved in cAMP signaling in cells.
Cellular Localization	Cell membrane
Post-translational Modifications	The pseudophosphorylation site binds to the substrate-binding region of the catalytic chain, resulting in the inhibition of its activity.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com