

Anti-PRIC285 antibody



Description	Rabbit polyclonal to PRIC285.
--------------------	-------------------------------

Model	STJ95222
Host	Rabbit
Reactivity	Human
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human PRIC285
Immunogen Region	1880-1960 aa, Internal
Gene ID	85441
Gene Symbol	HELZ2
Dilution range	WB 1:500-1:2000ELISA 1:20000
Specificity	PRIC285 Polyclonal Antibody detects endogenous levels of PRIC285 protein.
Tissue Specificity	Expressed in various tissues including heart, pancreas, skeletal muscle, colon, spleen, liver, kidney, lung, peripheral blood and placenta.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Helicase with zinc finger domain 2 ATP-dependent helicase PRIC285 Helicase with zinc finger 2, transcriptional coactivator PPAR-alpha-interacting complex protein 285 PPAR-gamma DNA-binding domain-interacting protein 1 PDIP1 <

Molecular Weight	300 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:30021 OMIM:611265
Alternative Names	Helicase with zinc finger domain 2 ATP-dependent helicase PRIC285 Helicase with zinc finger 2, transcriptional coactivator PPAR-alpha-interacting complex protein 285 PPAR-gamma DNA-binding domain-interacting protein 1 PDIP1 <
Function	Helicase that acts as a transcriptional coactivator for a number of nuclear receptors including PPARA, PPARG, THRA, THRB and RXRA.
Sequence and Domain Family	Contains 5 Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. These motifs are not required for interaction with PPARG.
Cellular Localization	Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com